

## **2008 Mathematics**

# Intermediate 2 – Units 1, 2 and 3 Paper 1

## **Finalised Marking Instructions**

#### © The Scottish Qualifications Authority 2008

The information in this publication may be reproduced to support SQA qualifications only on a non-commercial basis. If it is to be used for any other purposes written permission must be obtained from the Assessment Materials Team, Dalkeith.

Where the publication includes materials from sources other than SQA (secondary copyright), this material should only be reproduced for the purposes of examination or assessment. If it needs to be reproduced for any other purpose it is the centre's responsibility to obtain the necessary copyright clearance. SQA's Assessment Materials Team at Dalkeith may be able to direct you to the secondary sources.

These Marking Instructions have been prepared by Examination Teams for use by SQA Appointed Markers when marking External Course Assessments. This publication must not be reproduced for commercial or trade purposes.

### **General Marking Principles**

These principles describe the approach to be taken when marking Intermediate 2 Mathematics papers. For more detailed guidance please refer to the notes which are included with the Marking Instructions.

- 1 Marks must be assigned in accordance with the Marking Instructions. The main principle in marking scripts is to give credit for the skills demonstrated and the criteria met. Failure to have the correct method may not preclude a candidate gaining credit for the calculations involved or for the communication of the answer.
- 2 The answer to one part of a question, even if incorrect, must be accepted as a basis for subsequent dependent parts of the question. Full marks in the dependent part(s) may be awarded provided the question is not simplified.
- **3** The following should not be penalised:
  - working subsequent to a correct answer (unless it provides firm evidence that the requirements of the question have not been met)
  - omission or misuse of units (unless marks have been specifically allocated for the purpose in the marking scheme)
  - bad form, eg sin  $x^\circ = 0.5 = 30^\circ$
  - legitimate variation in numerical values / algebraic expressions.
- 4 Solutions which seem unlikely to include anything of relevance must nevertheless be followed through. Candidates still have the opportunity of gaining one mark or more provided the solution satisfies the criteria for the mark(s).
- 5 Full credit should only be given where the solution contains appropriate working. Where the correct answer may be obtained by inspection or mentally, credit may be given, but reference to this will be made in the Marking Instructions.
- 6 In general markers will only be able to give credit for answers if working is shown. A wrong answer without working receives no credit unless specifically mentioned in the Marking Instructions. The rubric on the outside of the question papers emphasises that working must be shown.
- 7 Sometimes the method to be used in a particular question is explicitly stated; no credit should be given where a candidate obtains the correct answer by an alternative method.
- 8 Where the method to be used in a particular question is not explicitly stated, full credit must be given for alternative methods which produce the correct answer.
- 9 Do not penalise the same error twice in the same question.
- 10 Do not penalise a transcription error unless the question has been simplified as a result.
- 11 Do not penalise inadvertent use of radians in trigonometry questions, provided their use is consistent within the question.

### **Practical Details**

The Marking Instructions should be regarded as a working document and have been developed and expanded on the basis of candidates' responses to a particular paper. While the guiding principles of assessment remain constant, details can change depending on the content of a particular examination paper in a given year.

- 1 Each mark awarded in a question is referenced to one criterion in the marking scheme by means of a bullet point.
- 2 Where a candidate has scored zero marks for any question attempted, "0" should be shown against the answer in the place in the margin.
- 3 Where a marker wishes to indicate how s/he has awarded marks, the following should be used:
  - (a) Correct working should be ticked,  $\checkmark$ .
  - (b) Where working subsequent to an error is followed through, if otherwise correct and can be awarded marks, it should be marked with a crossed tick, X.
  - (c) Each error should be underlined at the point in the working where it first occurs.

#### 4 Do not write any comments, words or acronyms on the scripts.

Question No	Marking Scheme Give 1 mark for each •	Illustrations of evidence for awarding a mark at each •
1	Ans: gradient is 4	
	• <sup>1</sup> interpret: find gradient	• <sup>1</sup> 4
		1 mark
NOTES:		
1. F	For an answer of $m = 4$ , $c = 5$	award 0/1
2. F	For 4 <i>x</i>	award 0/1
2	Ans: $3x^2 - 5x - 10$	
	• <sup>1</sup> process: start to multiply out brackets	• <sup>1</sup> evidence of 2 correct terms (eg $3x^2 - 15x$ )
	• <sup>2</sup> process: complete process of multiplying out brackets	• <sup>2</sup> $3x^2 - 15x + 2x - 10$
	• <sup>3</sup> process: collect like terms which must include $x^2$ term	$\bullet^3 \qquad 3x^2 - 5x - 10$
		3 marks
NOTES:	·	

### Mathematics Intermediate 2: Paper 1, Units 1, 2 and 3 (non-calc)

Question No	Marking Scheme Give 1 mark for each •	Illustrations of evidence for awarding a mark at each •
3 (a)	<b>Ans: 12th</b> • <sup>1</sup> interpret: interpret diagram	• <sup>1</sup> 12th <b>1 mark</b>
NOTES:		
(b)	Ans: 5/20 or equivalent • <sup>1</sup> process: calculate probability	$\bullet^1  \frac{5}{20}$ 1 mark
<b>NOTES:</b> 1. 4	Accept variations eg 5 : 20 0·25 25% 5 out of 20	

QuestionMarking SchemeNoGive 1 mark for each •		Illustrations of evidence for awarding a mark at each •		
4 (a)	Ans: $(x+y)(x-y)$			
	• <sup>1</sup> process: factorise correctly	• <sup>1</sup> $(x+y)(x-y)$		
		1 mark		
NOTES:				
(b)	Ans: 86			
	• <sup>1</sup> strategy: know to substitute in expression	• <sup>1</sup> $(9 \cdot 3 + 0 \cdot 7)(9 \cdot 3 - 0 \cdot 7)$		
	• <sup>2</sup> process: evaluate expression	• <sup>2</sup> 86		
		2 marks		
NOTES:	·			
1. <u>/</u>	Alternative method			
•	<sup>1</sup> strategy: know how to evaluate expression	• vidence of $9.3 \times 9.3$ $-0.7 \times 0.7$		
•	$\bullet^2$ process: evaluate expression	• <sup>2</sup> 86		
2. 1	For $9 \cdot 3^2 - 0 \cdot 7^2$ = $81 \cdot 9 - 4 \cdot 9$ = 77, with no additional working,	award 0/2		

Question No	Marking Scheme Give 1 mark for each •	Illustrations of evidence for awarding a mark at each •
5 (a)	<ul> <li>Ans: 1, 3, 6, 11, 16, 22, 24, 25</li> <li>•<sup>1</sup> communicate: table with cumulative frequency column</li> </ul>	• <sup>1</sup> 1, 3, 6, 11, 16, 22, 24, 25 <b>1 mark</b>
NOTES:		
(b)	Ans: $Q_2 = 4, Q_1 = 2 \cdot 5, Q_3 = 5$	
	• <sup>1</sup> communicate: state median	• <sup>1</sup> $Q_2 = 4$
	• <sup>2</sup> communicate: state lower quartile	• $Q_2 = 4$ • $Q_1 = 2 \cdot 5$ • $Q_3 = 5$
	• <sup>3</sup> communicate: state upper quartile	• <sup>3</sup> $Q_3 = 5$
		3 marks
NOTES:		
V	Where the quartiles have been obtained from	
(	(i) Number of books leading to $Q_2 = 3.5, Q_1 = 1.5, Q_3 = 5.5$	award 0/3
(	(ii) Frequency (unordered) leading to $Q_2 = 5, Q_1 = 2.5, Q_3 = 4$	award 0/3
(	(iii) Frequency (ordered) leading to $Q_2 = 2.5, Q_1 = 1.5, Q_3 = 5$	award 0/3
(	(iv) Cumulative frequency leading to $Q_2 = 13.5, Q_1 = 4.5, Q_3 = 23$	award 0/3
(c)	Ans: 1.25	
	• <sup>1</sup> process: calculate SIQR	• <sup>1</sup> 1.25
		1 mark
NOTES:	1	- 1

Question No	Marking Scheme Give 1 mark for each •	Illustrations of evidence for awarding a mark at each •
(d)	Ans: number of textbooks more spread out for girls	1
	• <sup>1</sup> communicate: a valid statement	• <sup>1</sup> a valid statement <b>1 mark</b>
NOTES:		
6	Ans: 40 sq cm	
	• <sup>1</sup> strategy: know how to find area	• <sup>1</sup> area = $\frac{1}{2} \times 16 \times 20 \times \frac{1}{4}$
	• <sup>2</sup> process: calculate area correctly	• <sup>2</sup> 40
		2 marks
NOTES:		
1. F	For $(\frac{1}{2} \times 16 \times 20 \times \sin \frac{1}{4})$ leading to an answer of	$140 \text{ cm}^2$ award $1/2$
2. I	For an answer of 40 cm <sup>2</sup> , without working	award1/2
3. I	For an answer of 160 cm <sup>2</sup> ( $\frac{1}{2} \times 16 \times 20$ )	award 0/2

Question No	Marking Scheme Give 1 mark for each •	Illustrations of evidence for awarding a mark at each •	
7	Ans: 19°		
	• <sup>1</sup> process: state the size of $\angle ABD$	• <sup>1</sup> 90°	
	• <sup>2</sup> process: calculate the size of $\angle BAD$	$\bullet^2$ 44°	
	• <sup>3</sup> process: calculate the size of $\angle BAC$	• <sup>3</sup> 19°	
		3 marks	
NOTES:			
1. <u>4</u>	Alternative method		
•	• <sup>2</sup> process: calculate the size of $\angle$ BEA (where E is the point of intersection of AC and BD) • <sup>2</sup> 71°		
	Angle ABD, angle BAD and angle BEA may not be a diagram and can be awarded the first and second r		
3. 4	A correct answer, without working.	award 3/3	
8	Ans: $a = 5, b = 3$		
	• <sup>1</sup> communicate: state the value of a	• <sup>1</sup> 5	
	• <sup>2</sup> communicate: state the value of b	• <sup>2</sup> 3	
		2 marks	
<b>NOTES:</b> 1. H	For $a = 3, b = 5$	award 1/2	

Question No	Marking Scheme Give 1 mark for each ●		Illustrations of evidence for awarding a mark at each •		
9 (a)	Ans: $a = -5, b = 3$	l			
	• <sup>1</sup> communicate:	state value of a	$\bullet^1$	-5	
	• <sup>2</sup> communicate:	state value of b	•2	1	
					2 marks
NOTES:					
(b)	Ans: $x = 5$				
	• <sup>1</sup> communicate:	correctly state equation of axis of symmetry	•1	<i>x</i> = 5	1 mark
NOTES:					
1. F	For any answer other	than $x = 5$			award 0/1
(c)	Ans: P(0, 26), Q(	(10, 26)			
	• <sup>1</sup> communicate:	state <i>x</i> -coordinates of P and Q	•1	(0, ?) and (10, ?)	
	• <sup>2</sup> strategy:	know how to find <i>y</i> -coordinate of P (or Q)	•2	$y = (0-5)^2 + 1$	
	$\bullet^3$ process:	find coordinates of P and Q	•3	P (0, 26), Q (10, 26)	
					3 marks
NOTES:	<u> </u>		<u>I</u>		

- 1. Where a candidate substitutes both 0 and 10 into an incorrect equation leading to different y coordinates for P and Q, all 3 marks are available
- 2. The third mark is available only when the y coordinates have been obtained by substitution into the quadratic equation

Question No	Marking Scheme Give 1 mark for each •	Illustrations of evidence for awardin a mark at each •	
10	Ans: $\frac{4}{3}$		
	• <sup>1</sup> strategy: know to use $\sin x / \cos x = \tan x$	$\bullet^1  \tan x = 4/5 \div 3/5$	
	• <sup>2</sup> process: calculate tan correctly	$\bullet^2  \frac{4}{3}$	
		2 marks	

TOTAL MARKS FOR PAPER 1 30

### [END OF MARKING INSTRUCTIONS]



## **2008 Mathematics**

# Intermediate 2 – Units 1, 2 and 3 Paper 2

# **Finalised Marking Instructions**

© The Scottish Qualifications Authority 2008

The information in this publication may be reproduced to support SQA qualifications only on a non-commercial basis. If it is to be used for any other purposes written permission must be obtained from the Assessment Materials Team, Dalkeith.

Where the publication includes materials from sources other than SQA (secondary copyright), this material should only be reproduced for the purposes of examination or assessment. If it needs to be reproduced for any other purpose it is the centre's responsibility to obtain the necessary copyright clearance. SQA's Assessment Materials Team at Dalkeith may be able to direct you to the secondary sources.

These Marking Instructions have been prepared by Examination Teams for use by SQA Appointed Markers when marking External Course Assessments. This publication must not be reproduced for commercial or trade purposes.

These principles describe the approach to be taken when marking Intermediate 2 Mathematics papers. For more detailed guidance please refer to the notes which are included with the Marking Instructions.

- 1 Marks must be assigned in accordance with the Marking Instructions. The main principle in marking scripts is to give credit for the skills demonstrated and the criteria met. Failure to have the correct method may not preclude a candidate gaining credit for the calculations involved or for the communication of the answer.
- 2 The answer to one part of a question, even if incorrect, must be accepted as a basis for subsequent dependent parts of the question. Full marks in the dependent part(s) may be awarded provided the question is not simplified.
- **3** The following should not be penalised:
  - working subsequent to a correct answer (unless it provides firm evidence that the requirements of the question have not been met)
  - omission or misuse of units (unless marks have been specifically allocated for the purpose in the marking scheme)
  - bad form, eg sin  $x^\circ = 0.5 = 30^\circ$
  - legitimate variation in numerical values / algebraic expressions.
- 4 Solutions which seem unlikely to include anything of relevance must nevertheless be followed through. Candidates still have the opportunity of gaining one mark or more provided the solution satisfies the criteria for the mark(s).
- 5 Full credit should only be given where the solution contains appropriate working. Where the correct answer may be obtained by inspection or mentally, credit may be given, but reference to this will be made in the Marking Instructions.
- 6 In general markers will only be able to give credit for answers if working is shown. A wrong answer without working receives no credit unless specifically mentioned in the Marking Instructions. The rubric on the outside of the question papers emphasises that working must be shown.
- 7 Sometimes the method to be used in a particular question is explicitly stated; no credit should be given where a candidate obtains the correct answer by an alternative method.
- 8 Where the method to be used in a particular question is not explicitly stated, full credit must be given for alternative methods which produce the correct answer.
- 9 Do not penalise the same error twice in the same question.
- 10 Do not penalise a transcription error unless the question has been simplified as a result.
- 11 Do not penalise inadvertent use of radians in trigonometry questions, provided their use is consistent within the question.

The Marking Instructions should be regarded as a working document and have been developed and expanded on the basis of candidates' responses to a particular paper. While the guiding principles of assessment remain constant, details can change depending on the content of a particular examination paper in a given year.

- 1 Each mark awarded in a question is referenced to one criterion in the marking scheme by means of a bullet point.
- 2 Where a candidate has scored zero marks for any question attempted, "0" should be shown against the answer in the place in the margin.
- 3 Where a marker wishes to indicate how s/he has awarded marks, the following should be used:
  - (a) Correct working should be ticked,  $\checkmark$ .
  - (b) Where working subsequent to an error is followed through, if otherwise correct and can be awarded marks, it should be marked with a crossed tick, X.
  - (c) Each error should be underlined at the point in the working where it first occurs.
- 4 Do not write any comments, words or acronyms on the scripts.

### Mathematics Intermediate 2: Paper 2, Units 1, 2 and 3

Question No	Marking Scheme Give 1 mark for each •	Illustrations of evidence for awarding a mark at each •	
1	Ans: £9625.93		
	• <sup>1</sup> strategy: know how to increase by $4.5\%$	$\bullet^1$ × 1.045	
	• <sup>2</sup> strategy: know how to calculate amount	• <sup>2</sup> 50 000 × 1.045 <sup>4</sup>	
	$\bullet^3$ strategy: know how to calculate interest	• <sup>3</sup> 50 000 × $1.045^4 - 50\ 000$	
	• <sup>4</sup> process: carry out all calculations correctly within a valid strategy and round to nearest penny	• <sup>4</sup> 9625.93	
		4 marks	
NOTES:			
1	For an answer of £9625.93, with or without working	g award 4/4	
2	For an answer of £59 625.93, with or without worki	award 3/4	
3	For an answer of £2567.62 (the fourth year's interes	st), with working award 3/4	
4	Where an incorrect percentage has been used, the w through to give the possibility of awarding $3/4$	orking must be followed	
	eg for an answer of £171 025.31 (50 000 × $1.45^4$ – 5	50 000), with working award 3/4	
5	For an answer of £41 589.48 (50 000 $\times$ 0.955 <sup>4</sup> ) award 2/4		
6	For an answer of £8410.52 (50 000 – 41 589.48) award 2/4		
7	For an answer of £9000 (50 000 $\times$ 0.045 $\times$ 4) award 0/4		

Question No	Marking Scheme         Give 1 mark for each ●         Ans: 58 600 cubic cm			Illustrations of evidence for awarding a mark at each •	
2 (a)					
	$\bullet^1$ strategy:	know how to calculate vol of basket	lume • <sup>1</sup>	volume of cuboid cylinder	+ volume of
	• <sup>2</sup> process:	substitute correctly into vo formulae	blume $\bullet^2$	$30 \times 24 \times 50 + \pi \times 1$	$2^2 \times 50$
	• <sup>3</sup> process:	calculate total volume	•3	58 619 cm <sup>3</sup>	
	• <sup>4</sup> process:	round answer to 3 signific figures	ant $\bullet^4$	58 600 cm <sup>3</sup>	
					4 marks
NOTES:					
1 4	Accept variation	ns in volume due to variations	s in the valu	e of $\pi$	
		t is available for rounding an er requires no rounding, the			nt figures.
3 (	Common wrong	answers			
	43 200	(cuboid + sphere)	with wo	-	award 3/4
	47 300	(cuboid + $\frac{1}{2}$ cylinder)	with wo	e	award 3/4
	39 800 1170	(cuboid + $\pi dh$ ) (area of cross section)	with wo with wo	÷	award 3/4 award 2/4
	36 500	(cuboid + $\pi r^2$ )	with wo	•	award 2/4 award 2/4
(b)	Ans: 29.9 cl	m			
	• <sup>1</sup> strategy:	know how to find expression volume of box	n for $\bullet^1$	$35 \times 28 \times h$	
	• <sup>2</sup> process:	equate volume with $\frac{1}{2}$ of an to part (a)	$\bullet^2$	$35 \times 28 \times h = \frac{1}{2} \times 54$	8 600
	• <sup>3</sup> communi	cate: state value for <i>h</i>	•3	<i>h</i> = 29.9	
					3 marks
	<u> </u>				
NOTES:					

Question No	Marking Scheme Give 1 mark for each ●	Illustrations of evidence for awarding a mark at each ●		
3 (a)	Ans: 14.8			
	• <sup>1</sup> process: calculate the mean	• <sup>1</sup> 27		
	• <sup>2</sup> process: calculate $(x - \overline{x})^2$	• <sup>2</sup> 289, 81, 1, 25, 484		
	• <sup>3</sup> process: substitute into formula	• <sup>3</sup> $\sqrt{(880/4)}$		
	• <sup>4</sup> process: calculate standard deviation	• <sup>4</sup> 14.8 (disregard rounding)		
		4 marks		
NOTES:				
1. <u>Alter</u>	mative method			
$\bullet^1$	process: calculate $\sum x$ and $\sum x^2$	• <sup>1</sup> 135 and 4525		
•2	process: substitute into formula	• <sup>2</sup> $\sqrt{\frac{4525 - 135^2 / 5}{5 - 1}}$		
•3	process: simplify	$\bullet^3  \sqrt{\frac{880}{4}}$		
•4	process: calculate standard deviation	• <sup>4</sup> 14.8 (disregard rounding)		
2 For c	correct answer, without working	award 0/4		
(b)	Ans: The physics marks were more consistent than the maths marks (since 6.8 < 14.8)			
	• <sup>1</sup> communicate: valid comment about the spread of marks	• <sup>1</sup> valid comment 1 mark		
NOTES:		1 m		

Question No	Marking Scheme Give 1 mark for each ●	Illustrations of evidence f a mark at each	0
(c)	<b>Ans:</b> $y = \frac{1}{2}x + 20$		
	• <sup>1</sup> process: find gradient	• <sup>1</sup> $m = \frac{1}{2}$ (or equivalent)	
	• <sup>2</sup> process: state <i>y</i> -intercept or $c$ in $y = mx + c$	$\bullet^2$ c = 20	
	• <sup>3</sup> communicate: state equation of line	$\bullet^3  y = \frac{1}{2}x + 20$	
			3 marks
NOTES:			
1 H	For correct answer without working		award 3/3
2 H	For $p = 0.5m + 20$		award 3/3
3 H	For $y = 0.5x$		award 1/3
	Where m and/or c are incorrect the working must b he possibility of awarding 1/3 or 2/3	e followed through to give	
	f the equation is stated incorrectly and there is no for correct gradient or correct <i>y</i> -intercept	working, 1/3 can be awarded	
	For an incorrect equation (ie both m and c incorrect equation $y = 20x + 0.5$	t), without working	award 0/3
(d)	Ans: 58%		
	• <sup>1</sup> process: calculate physics % using equation	•1 $y = \frac{1}{2}(76) + 20 = 58$	
			1 mark
NOTES:			1 ma

Question No	Marking Scheme Give 1 mark for each •	Illustrations of evidence for awarding a mark at each •
4 (a)	Ans: $280x + 70y = 5250$ • <sup>1</sup> interpret: interpret the text	• <sup>1</sup> $280x + 70y = 5250$ <b>1 mark</b>
<b>NOTES:</b> 1 <i>A</i>	$Accept \ 280x + 70y = 52.50$	
(b)	Ans: $210x + 40y = 3800$ • <sup>1</sup> interpret: interpret the text	• $^{1}$ 210x + 40y = 3800 1 mark
<b>NOTES:</b> 1 <i>A</i>	Accept $210x + 40y = 38.00$ when consistent with the	e answer to part (a)

Question No	Marking Scheme Give 1 mark for each •	Illustrations of evidence for awarding a mark at each •	
4 (c)	Ans: Calls cost 16 pence per minute, texts cost 11 pence each		
	• <sup>1</sup> strategy: know to solve system of equations	• <sup>1</sup> evidence	
	• <sup>2</sup> process: follow a valid strategy through to produce a value for $x$ and $y$	• <sup>2</sup> a value for x and y	
	• <sup>3</sup> process: correct value for $x$ and $y$	• <sup>3</sup> $x = 16, y = 11$	
	• <sup>4</sup> communicate: state result	• <sup>4</sup> a call costs 16p per minute a text costs 11 pence	
		4 marks	
NOTES:	L		
1 1	Incorrect equations must be followed through to give	te the possibility of awarding 4/4	
2	Any valid strategy must involve the use of two equa	tions	
6	Where the correct values for x and y have been obtain equations, marks are available only if both values have equations we $280 \times 16 + 70 \times 11 = 5250$ $210 \times 16 + 40 \times 11 = 3800$ leading to $x = 16$ , $y = 11$ a call costs 16p per minute a text costs 11p		
4 1	For $x = 16$ , $y = 11$ (with working) award 3/4 (loses of	communication mark)	
	For the award of the final mark the price of a call perstated in pence or pounds	er minute and the price of a text must be	
	For a wrong answer without working, or based on an invalid strategy, the final mark cannot be awarded		
7 1	For a correct answer, without working	award 0/4	

Question No	Marking Scheme Give 1 mark for each •	Illustrations of evidence for awarding a mark at each •
5	Ans: Angle EDF = 111.8°	
	• <sup>1</sup> strategy: know to apply cosine rule to find angle EDF	• <sup>1</sup> evidence
	• <sup>2</sup> process: correct application of cosine rule	• <sup>2</sup> $\cos D = \frac{10.4^2 + 13.2^2 - 19.6^2}{2 \times 10.4 \times 13.2}$
	• <sup>3</sup> process: calculate angle EDF	• <sup>3</sup> 111.8°
		3 marks
NOTES:		
	Where an angle other than angle EDF has been calc maximum of 2/3 can be awarded provided that the v with the application of the cos rule	
2	1.95 (RAD), 124.2 (GRAD), with working	award 3/3
3	For an answer obtained by scale drawing,	award 0/3

Question No	Marking Scheme Give 1 mark for each •	Illustrations of evidence for awarding a mark at each •
6	Ans: 0.35, – 1.15	
	• <sup>1</sup> strategy: know to use quadratic formula	• <sup>1</sup> evidence
	• <sup>2</sup> process: correct substitution in formula	• <sup>2</sup> $\frac{-4 \pm \sqrt{(4)^2 - 4(5)(-2)}}{2(5)}$
	• <sup>3</sup> process: calculate $b^2 - 4ac$ correctly	• <sup>3</sup> 56
	• <sup>4</sup> process: state both values of $x$ correct to two decimal places	•4 0.35, -1.15
		4 marks
NOTES:		2
	Where $b^2 - 4ac$ is calculated incorrectly, the fourth	mark is available only if $b^2 - 4ac > 0$
2	Alternative method (graphical solution)	
•	• strategy: know to graph $y = 5x^2 + 4x - 2$	• <sup>1</sup> $\bigstar \mathcal{Y}$
		$y = 5x^2 + 4x - 2$
	• <sup>2</sup> communicate: indicate position of roots	• <sup>2</sup> $y = 5x^2 + 4x - 2$

- $\bullet^3$  communicate: state first root correct to 2 •<sup>3</sup> -1.15decimal places
- communicate: state second root correct to 2 •4 •<sup>4</sup> 0·35 decimal places
- For a correct answer, without working 3

award 0/4

Question No	Marking Scheme Give 1 mark for each •	Illustrations of evidence for awarding a mark at each •
7 (a)	Ans: $m^2$	
	process: simplify indices	$\bullet^1$ $m^2$
		1 mark
NOTES:		
(b)	Ans: $\sqrt{5}$	
	• <sup>1</sup> process: simplify surd $\sqrt{20}$	$\bullet^1$ $2\sqrt{5}$
	• <sup>2</sup> process: simplify surd $\sqrt{45}$	$\bullet^2$ $3\sqrt{5}$
	• <sup>3</sup> process: state answer in simplest form	$\bullet^3$ $\sqrt{5}$
		3 marks
NOTES:		
1 F	For correct answer, without working	award 0/3
8	Ans: $x = 138.6, 221.4$	
	• <sup>1</sup> process: start to solve equation	• <sup>1</sup> $\cos x^\circ = -3/4$
	• <sup>2</sup> process: calculate one value of $x$	• <sup>2</sup> 138.6
	• <sup>3</sup> process: calculate second value of $x$	• <sup>3</sup> 221.4
		3 marks
NOTES:		
	Where $\cos x^{\circ} > 0$ , 1/3 can be awarded when 2 values ncorrect value for $\cos x^{\circ}$ (working eased)	s of $x$ are calculated consistent with the
	Where a graphical solution has been used, the first r what graph is drawn and where the values occur	nark is available for indicating
3 F	For correct answer, without working	award 0/3

Question No	Marking Scheme Give 1 mark for each •	Illustrations of evidence for awarding a mark at each •
9	Ans: 16 cm	
	• <sup>1</sup> strategy: marshall facts and know to use right-angled triangle	•1 10
	• <sup>2</sup> strategy: know that PQ bisects AB	• <sup>2</sup> 10 6
	• <sup>3</sup> process: use Pythagoras' Theorem	• <sup>3</sup> $x^2 = 10^2 - 6^2$
	• <sup>4</sup> process: calculate length of third side	• <sup>4</sup> $x = 8$
	• <sup>5</sup> process: calculate PQ	• <sup>5</sup> 16 cm <b>5 marks</b>
NOTES:	1	1
I	SPECIAL CASE: Where $\angle PAQ = 90^{\circ}$ or $\angle APQ = \angle AQP = 45^{\circ}$ are available for correct Pythagoras or Trigonometric ca	
2 8	SOME COMMON ANSWERS (with working)	
<u>/</u>	Answer	Maximum mark available
,	$2 \times \sqrt{10^2 + 6^2} = 23 \cdot 32$	4/5
	$\sqrt{10^2 + 6^2} = 11.66$	3/5
2	$2 \times \sqrt{12^2 - 10^2} = 13 \cdot 27$	3/5
	$\sqrt{12^2 - 10^2} = 6 \cdot 63$	2/5
	$\sqrt{12^2 + 10^2} = 15 \cdot 62$	2/5
	$\sqrt{10^2 + 10^2} = 14 \cdot 14$	2/5 (see note 1)
3 H	For a correct answer, without working	award 0/5

Question No	Marking Scheme	Illustrations of evidence for awarding
INU	Give 1 mark for each •	a mark at each •
10	Ans: $(p-q)^2$	
	• <sup>1</sup> process: start to re-arrange formula	• <sup>1</sup> $\sqrt{a} = p - q$ • <sup>2</sup> $a = (p - q)^2$
	• <sup>2</sup> process: make <i>a</i> the subject	$\bullet^2  a = (p - q)^2$
		2 marks
NOTES:		
1 F	For a correct answer, with or without working	award 2/2
11	Ans: $\frac{8-a}{a(a+4)}$	
	• <sup>1</sup> process: state a valid common denominator	• <sup>1</sup> any valid denominator
	• <sup>2</sup> process: find correct numerator of equivalent fraction	$\bullet^2$ both numerators correct
	• <sup>3</sup> process: state answer in simplest form	$\bullet^3  \frac{8-a}{a(a+4)}$
		3 marks
NOTES:	1	1

TOTAL MARKS FOR PAPER 2 50

### [END OF MARKING INSTRUCTIONS]